

****REPRESENTATIVE DATASHEET******Goat anti-human von Willebrand Factor**

Whole IgG from antiserum

2.5 mg

Product #: GAVWF-IG-ASR**Lot #:** XXXX**Expiry date:** XXXX**DOM:** XXXX

Store at -10 to -20°C.

1395 Sandhill Drive. Ancaster, Ontario, Canada L9G 4V5
905-304-9896 • 800-903-6020 • fax 905-304-9897**Analyte Specific Reagent.**

Analytical and performance characteristics are not established.

Description of von Willebrand Factor

von Willebrand Factor (vWF, also previously referred to as Factor VIII related antigen) is a large adhesive protein produced in endothelial cells and megakaryocytes. There are two critical functions of vWF, the first being its involvement in the process of platelet adhesion and aggregation through interaction with platelet receptor glycoprotein Ib, the second being the binding and stabilization of Factor VIII (antihemophilic factor) for secretion and transport in plasma. The vWF precursor protein is synthesized with a 95,000 dalton propeptide (also known as vWF antigen-II), believed to be involved in the intracellular multimerization of the vWF subunits. The mature vWF multimers are then packed into storage organelles within the cell (Weibel-Palade bodies) after which the propeptide is cleaved and released. vWF circulates as multimers of disulphide linked 220,000 dalton subunits and the molecular weight of these multimers ranges from 0.5-20 million daltons.

The plasma concentration of vWF is typically 10 µg/ml, but increased levels are often observed in pregnancy and other conditions of physiological stress. von Willebrand's disease (vWD) is perhaps the most common inherited bleeding disorder in humans and is the result of either quantitative deficiencies of vWF (vWD Types I & III), or one of a number of qualitative disorders of vWF structure and function (vWD Type II).

REVIEW ARTICLES

1. Montgomery RR, Collier BS, in **Hemostasis and Thrombosis**, 3rd Edition, eds. RW Colman, J Hirsh, VJ Marder and EW Salzman, pp. 134-168, J.B. Lippincott Co., Philadelphia PA, USA, 1994.

2. Sadler JE; von Willebrand Factor (Minireview); JBC 266:34, pp 22777-22780, 1991.

Product Specifications**Description:**

Vial containing XXXX ml of whole IgG representing approximately 1 ml of antiserum. Total protein is 2.5 mg.

Format:

Whole IgG, clear liquid.

Host Animal:

Goat

Immunogen:

Human vWF purified from plasma.

Concentration:IgG concentration is XXXX mg/ml, determined by absorbance using an extinction coefficient ($E_{1\%}^{1\text{cm}}$) of 13.4.**Buffer:**

10 mM HEPES, pH 7.4, 150 mM NaCl, 50% (v/v) glycerol.

Storage:

Store between -10 and -20°C. Product will become viscous but will not freeze. Avoid storage in frost-free freezers. Keep vial tightly capped. Allow product to warm to room temperature and gently mix before use.

Specificity:

This antibody is specific for vWF as demonstrated by immunoelectrophoresis and ELISA.

Precautions:

Unused solution should be disposed of according to current local, State and Federal Regulations. For a Material Safety Data Sheet for this product contact Affinity Biologicals Inc.

Visit our site (www.affinitybiologicals.com) for other related products.

Limited Warranty: This product is warranted to perform in accordance with its labeling and literature. Affinity Biologicals Inc. disclaims any implied warranty of merchantability or fitness for any other purposes, and in no event will Affinity Biologicals Inc. be liable for any consequential damages arising out of aforesaid express warranty.

Manufactured in Canada by:
AFFINITY BIOLOGICALS INC.
1395 Sandhill Drive
Ancaster ON CANADA L9G 4V5
Tel: (905) 304-9896
(800) 903-6020
Fax: (905) 304-9897
info@affinitybiologicals.com